Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at http://about.jstor.org/participate-jstor/individuals/early-journal-content.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

# RATIONAL EDGED CUBOIDS WITH EQUAL VOLUMES AND EQUAL SURFACES.

By L. E. DICKSON, The University of Chicago.

1. In the BULLETIN, May, 1909, p. 401, Professor Kasner proposed the problem to find two cuboids (rectangular parallelopipeds) with equal volumes and equal surfaces, and in which the dimensions are all integral.

The problem is to find two distinct triples of integers such that

$$xyz=x'y'z', \quad xy+xz+yz=x'y'+x'z'+y'z'. \tag{1}$$

I shall prove that each integer must exceed unity and that the volume $xyz$ must be the product of five or more primes. Among the solutions which I obtain, the simplest are

$$p,\ p,\ \tfrac{1}{2}(p+1)p^2; \quad p^2,\ p^2,\ \tfrac{1}{2}(p+1) \qquad (p \text{ odd}), \tag{2}$$

$$p,\ (p+1)q,\ (p+1)(p+1-q); \quad p+1,\ (p+1)p,\ q(p+1-q) \quad (1<q<p), \tag{3}$$

in which $q=r$ and $q=p+1-r$ give the same sets. While for (3) the volume contains at least six primes, for (2) there may be only five primes, for example,

(2′) 3, 3, 18; 9, 9, 2. 5, 5, 75, 25, 25, 3.

The simplest examples under (3) are

(3′) 3, 8, 8; 4, 12, 4. 4, 10, 15; 5, 20, 6.

Generalizations of (2) and (3) are given by (9)-(10) and (16)-(17).

The only solutions in which $xyz<200$ are the first solutions under (2′) and (3′).

In §§7-9, I show that for three rational edged cuboids with equal volumes and equal surfaces, the volume is the product of six or more primes, that no two of the triples are of the type $p_1,\ p_2,\ p_3p_4p_5p_6$, and that the cases in which two of the triples are $p_1,\ p_2,\ p_3p_4p_5p_6$ and $p_3,\ p_1p_2,\ p_4p_5p_6$ are excluded.

2. Theorem. *Two triples are identical if they have an element in common.*

If $x=x'$, equations (1) reduce to $yz=y'z'$, $y+z=y'+z$.

3. Theorem. *In any solution each integer exceeds unity.*

Two sets with equal products and having an element unity must be of the form

$$(4) \qquad 1,\ f_1f_2f_3,\ f_4f_5f_6;\ f_1f_4,\ f_2f_5,\ f_3f_6.$$

Denote the corresponding surfaces by $2S_1$ and $2S_2$. Then

$$S_1=f_1f_2f_3+f_4f_5f_6+f_1\ldots f_6, \qquad S_2=f_1f_2f_4f_5+f_1f_3f_4f_6+f_2f_3f_5f_6.$$

First, let each $f_i>1$. Since the substitution (12)(45) leaves the triples (4) unaltered, we may set $f_4 \geqq f_5$.

Now $(f_i-1)(f_j-1) \geqq 1$, so that $f_if_j \geqq f_i+f_j$. Thus

$$f_1\ldots f_6 \geqq f_1f_2f_4f_6(f_3+f_5) \geqq f_3f_4f_6(f_1+f_2)+f_1f_2f_5(f_4+f_6).$$

Also $f_3f_4f_6f_2 \geqq f_2f_3f_5f_6$. Hence $f_1\ldots f_6>S_2$, $S_1>S_2$.

Second, let a single $f_i$ be unity. We may set $f_3=1$, $f_5 \geqq f_4$. Then

$$f_1\ldots f_6 \geqq f_1f_2f_5(f_4+f_6) \geqq f_1f_2f_4f_5+f_5f_6(f_1+f_2) \geqq S_2,$$

since $f_1f_5f_6 \geqq f_1f_4f_6$. Hence $S_1>S_2$.

Third, let at least two $f_i$ equal unity. If two such $f$'s belong to the same product in $(4_1)$, we may set $f_2=f_3=1$. Then

$$S_1=f_1+f_4f_5f_6+f_1f_4f_5f_6>f_5f_6+f_1f_4(f_5+f_6), \quad S_1>S_2,$$

if $f_5>1$, $f_6>1$. If $f_5=1$ or $f_6=1$, the triples have the common element 1 and are identical by §2. Next, let the two $f$'s which equal unity belong to different products in $(4_1)$. A case like $f_3=f_6=1$ is excluded by §2. Hence we may set $f_3=1$, $f_4=1$. Thus

$$S_1-S_2=f_2f_5(f_1f_6-f_1-f_6)+f_1f_2+f_5f_6-f_1f_6.$$

If $f_2>1$ or $f_5>1$, $S_1-S_2>(f_2f_5-1)(f_1f_6-f_1-f_6)$, so that $S_1-S_2>0$ unless $f_1=1$ or $f_6=1$. In the latter case the triples (4) have a common element 1. The same is true in the remaining case $f_2=f_5=1$.

4. Theorem. *The volume must contain at least five prime factors.*

By §3, each element exceeds 1. Hence there occur four or more primes. If there are just four primes, the triples without a common element (§2) may be designated

$$(5) \qquad p_1,\ p_2,\ p_3p_4;\ \ p_3,\ p_4,\ p_1p_2,$$

$p_4$ being the greatest $p$, and $p_2 \geqq p_1$. Then $S_1+p_1p_2p_3p_4=S_2+p_1p_2p_3p_4$ may be written

$$p_1p_2(p_3-1)(p_4-1)=p_3p_4(p_1-1)(p_2-1).$$

Hence $p_4=p_2$, so that the triples (5) are identical by §2.

5. For five primes, each triple must be of the form (1, 1, 3) or (1, 2, 2), the notation indicating the number of prime factors of each element. Consider first

$$(6) \qquad p_1,\ p_2,\ p_3p_4p_5;\ \ p_3,\ p_1p_2,\ p_4p_5,$$

where $p_1 \geqq p_2$. From $S_1+p_1\ldots p_5=S_2+p_1\ldots p_5$, we get

$$p_1p_2(p_3-1)(p_4p_5-1)=p_3p_4p_5(p_1-1)(p_2-1).$$

Hence $p_1$ divides $p_3p_4p_5$. But $p_1 \neq p_3$ by §2. Since (6) is unaltered by the interchange of $p_4$ and $p_5$, we may set $p_1=p_4$. Thus

$$(7) \qquad p_2(p_3-1)(p_1p_5-1)=p_3p_5(p_1-1)(p_2-1) \qquad (p_1 \geqq p_2).$$

If $p_2$ does not divide $p_1-1$, then $p_2=p_5$, and (7) becomes

$$(8) \qquad p_3(p_1+p_2-2)=p_1p_2-1,\ \ (p_3-1)^2=(p_1-p_3)(p_2-p_3).$$

Let $g$ be the greatest common divisor of $p_1-p_3=ga^2$, $p_2-p_3=gb^2$. Then $p_3-1=gab$,

$$(9) \quad p_1=1+gab+ga^2,\ \ p_2=1+gab+gb^2,\ \ p_3=1+gab \ \ (a,\ b \text{ relatively prime})$$

*For these values we have the solution*

$$(10) \qquad p_1,\ p_2,\ p_1p_2p_3;\ \ p_3,\ p_1p_2,\ p_1p_2.$$

For $a=b$, then $a=1$, $p_3=\frac{1}{2}(p_1+1)$ and (10) becomes solution (2). For the remaining cases we may set $a>b$. Examples when the $p$'s are all primes are (2′) and

$a=2$, $b=1$, $g=2$ or 6, $p_1$, $p_2$, $p_3=13$, 7, 5 or 37, 19, 13;
$a=3$, $b=2$, $g=10$, $p_1=151$, $p_2=101$, $p_3=61$.

Next, let $p_2$ divide $p_1-1$. Then (7) is equivalent to

$$(11) \qquad p_1-1=cp_2,\ \ p_3-1=kp_5,\ \ p_1p_5-1=lp_3,\ \ lk=c(p_2-1).$$

From the second and third we eliminate $p_3$ and see that $l=-1+mp_5$, $m=p_1-lk$. Hence by the fourth and first, $m=1+c$. Hence $(11_3)$ may be replaced by

$$l=-1+(1+c)p_5. \tag{12}$$

Let $g$ be the greatest common divisor of $c$ and $k$. Then, by ($11_4$),

$$c=g\gamma,\ k=g\mu,\ l=\lambda\gamma,\ p_2-1=\lambda\mu \quad (\lambda \text{ and } \mu \text{ relatively prime}). \tag{13}$$

Set $\lambda-gp_5=\rho$. Then (12) becomes $\lambda\rho=p_5-1$. We may thus eliminate $p_5$ and $\lambda$:

$$p_5=1+\gamma\rho,\ p_2=1+\lambda\mu,\ p_1=1+g\gamma p_2,\ p_3=1+g\mu p_5 \quad (\lambda=g+\rho+g\rho\gamma). \tag{14}$$

*For any positive integers $g$, $\rho$, $\gamma$, $\mu$, of which the last two are relatively prime, formulae* (14) *give values of the $p_i$ leading to a solution**

$$p_1,\ p_2,\ p_1p_3p_5;\ p_3,\ p_1p_2,\ p_1p_5. \tag{15}$$

For $\gamma=\mu=1$, $p_2=(1+g)(1+\rho)$ is composite. For $\gamma=1$, $\mu=2$, $\rho=1$, the least value of $g$ giving prime $p$'s is $g=10$ and $p_5=2$, $p_2=43$, $p_1=431$, $p_3=41$.

6. For two triples of type (1, 2, 2), it suffices to consider

$$p_1,\ p_2p_3,\ p_4p_5;\ p_2,\ p_1p_4,\ p_3p_5 \quad (p_2>p_1;\ p_5\neq p_1,\ p_2;\ p_3\neq p_4). \tag{16}$$

Then $S_1=S_2$ gives $p_2p_3p_5(p_4-1)\equiv 0 \pmod{p_1}$. If $p_1=p_3$, $S_1=S_2$ gives

$$p_1p_2+p_4p_5+p_2p_4p_5=p_2p_4+p\ \ p_5+p_1p_4p_5.$$

Thus $p_4p_5(p_1-1)\equiv 0 \pmod{p_2}$, so that $p_2=p_4$. The middle terms in (16) would then be equal. This case is thus excluded by §2. Hence $p_4\equiv 1 \pmod{p_1}$.

Next, $S_1=S_2$ gives $p_1p_4p_5(p_3-1)\equiv 0 \pmod{p_2}$. First, let $p_3-1$ be prime to $p_2$, so that $p_4=p_2$. Removing the factor $p_2$ from $S_1=S_2$, we get

$$p_1(p_2-p_3-p_5)=p_3p_5(p_2-p_1-1).$$

Since $p_2\equiv 1 \pmod{p_1}$, $p_2-p_1-1=cp_1$, where $c$ is an integer $\geqq 0$. Thus

$$p_4=p_2=(c+1)p_1+1=p_3+p_5+cp_3p_5. \tag{17}$$

For $c=0$, we set $p_1=p$, $p_3=q$ and obtain solution (3); note that $p$ occurs in the second triple if and only if $q=1$ or $p$.

---

*To show that these triples are distinct, it suffices to prove that $p_1$ does not occur in the second. But $p_1<p_1p_2$, $p_1<p_1p_5$. If $p_1=p_3$, then $\gamma p_2=\mu p_5$. But $\gamma$ and $\mu$ are relatively prime, and $\gamma$ does not divide $p_5$ by ($14_1$).

For $c=1$, $2(p_1+1)=(p_3+1)(p_5+1)$. If $p_3=2$, then $p_1=3k-1$, $p_5=2k-1$. The lowest values of $k$ leading to prime values for each $p_i$ are $k=2, 4, 10$:

$$p_1,\ p_5,\ p_2=5,\ 3,\ 11;\ 11,\ 7,\ 23;\ 29,\ 19,\ 59.$$

The first gives the solution 5, 22, 33; 6, 11, 55. If $p_3$ is odd, then

$$p_3=2k-1,\quad p_1=k(p_5+1)-1,\quad p_4=p_2=2k(p_5+1)-1. \tag{18}$$

The $p_i$ are all primes for $k=2$, $p_5=5, 11$; $k=3$, $p_5=7, 13$:

$$p_5,\ p_3,\ p_1,\ p_2=5,\ 3,\ 11,\ 23;\ 11,\ 3,\ 23,\ 47;\ 7,\ 5,\ 23,\ 47;\ 13,\ 3,\ 41,\ 83.$$

For $c=2$, we have $p_3=3$, $p_5=3k+1$, $p_1=7k+3$, or

$$p_3=3l+1,\quad p_1=l+p_5+2lp_5,\quad \text{or}\quad p_3=3l+2,\quad p_5=3m+1,\quad p_1=6lm+3l+5m+2.$$

But for $c$ even, the $p$'s in (17) are not all primes if $p_1\neq 2$.

In view of the variety of solutions obtained, further cases are not considered.

7. Do there exist three distinct cuboids with rational edges having equal volumes and equal surfaces? We show that there is no solution in which the volume is the product of fewer than six primes. In view of §4, we consider the case of three triples of five primes $p_i$. Not all three are of the type (1, 1, 3). Let two be of type (1, 1, 3) and one of type (1, 2, 2); it suffices to consider

$$p_1,\ p_2,\ p_3p_4p_5;\ p_3,\ p_4,\ p_1p_2p_5;\ p_5,\ p_1p_2,\ p_3p_4,\ \text{or}\ p_5,\ p_1p_3,\ p_2p_4. \tag{19}$$

In the first alternative, we reduce the $S_i$ modulo $p_1$ and get

$$p_2p_3p_4p_5\equiv p_3p_4\equiv p_3p_4p_5 \pmod{p_1}.$$

Since $p_3p_5$ is not divisible by $p_1$, $p_5\equiv 1$, $p_2\equiv 1 \pmod{p_1}$. Similarly $p_1\equiv 1 \pmod{p_2}$, contrary to the former. Likewise ($19_2$) gives $p_3\equiv 1 \pmod{p_1}$, $p_1\equiv 1 \pmod{p_3}$ and is excluded.

Next, let one triple be of the type (1, 1, 3) and two of the type (1, 2, 2); it suffices to treat

$$(20) \qquad (p_1,\ p_2p_3,\ p_4p_5;\ p_2,\ p_1p_4,\ p_3p_5;\ p_3,\ p_4,\ p_1p_2p_5,\ \text{or}\ p_4,\ p_5,\ p_1p_2p_3.$$

But, in either case, we find that $p_4 \equiv 1 \pmod{p_1}$, $p_1 \equiv 1 \pmod{p_4}$.

Hence all three triples must be of type (1, 2, 2). Two of them may be taken to be the first two in (20), with $p_3 \neq p_4$, $p_2 \neq p_5$, $p_1 \neq p_5$. Since these are unaltered by (12) (34), the third may be restricted to one of the three:

$$p_3,\ p_1p_5,\ p_2p_4;\ p_5,\ p_1p_2,\ p_3p_4;\ p_5,\ p_1p_3,\ p_2p_4.$$

The first case gives $p_3 \equiv 1 \pmod{p_2}$, $p_2 \equiv 1 \pmod{p_3}$. The second case gives $p_2 \equiv 1 \pmod{p_1}$, $p_1 \equiv 1 \pmod{p_2}$. In the third case we may set $p_2 > p_1$; then

$$p_1p_4p_5 \equiv p_1p_2p_4 \equiv p_2p_4p_5 \pmod{p_3}, \quad p_1p_4p_5 \equiv p_1p_3p_4p_5 \equiv p_1p_3p_5 \pmod{p_2}.$$

If $p_3 = p_5$, then $p_1p_2p_4$ is a multiple of $p_3$, whereas $p_1 \neq p_5$, $p_2 \neq p_5$, $p_4 \neq p_3$. Hence $p_4p_5(p_2 - p_1) \equiv 0$, gives $p_2 - p_1 = cp_3$, $c > 0$. Similarly, $p_1p_4p_5(p_3 - 1) \equiv 0 \pmod{p_2}$ gives $p_3 \equiv 1 \pmod{p_2}$, since $p_4 = p_2$ would require $p_1p_3p_5 \equiv 0 \pmod{p_2}$. Hence $-p_1 \equiv c \pmod{p_2}$, so that $c \geqq p_2 - p_1$. The former equation $p_2 - p_1 = cp_3$ is thus impossible in view of $p_3 > 1$. *There is no set of three triples involving fewer than six primes.*

8. Consider three triples involving six primes. If all are of type (1, 1, 4), they may be taken to be

$$(21) \qquad p_1,\ p_2,\ p_3p_4p_5p_6;\ p_3,\ p_4,\ p_1p_2p_5p_6;\ p_5,\ p_6,\ p_1p_2p_3p_4,$$

with $p_3p_4 > p_5p_6$. By $S_2 = S_3$, $p_3p_4 = p_5p_6 + \delta \cdot p_1p_2$, $\delta > 0$. But $p_1p_2 \equiv p_5p_6 \pmod{p_3p_4}$ by $S_1 = S_3$. Hence $0 \equiv p_5p_6(1 + \delta) \pmod{p_3p_4}$. Thus $\delta = -1 + \varepsilon\, p_3p_4$, $\epsilon > 0$,

$$p_3p_4 + p_1p_2 = p_5p_6 + \varepsilon\, p_1p_2p_3p_4.$$

But $p_1p_3 \cdot p_2p_4 \geqq (p_1 + p_3)(p_2 + p_4) > p_1p_2 + p_3p_4$. Hence this case is excluded.

Let two of the triples be the first two in (21) and the third of type (1, 2, 3). Since the former are unaltered or interchanged by (12), (34), (56), (13) (24), the third may be assumed to be $p_5,\ p_1p_2,\ p_3p_4p_6$; or $p_5,\ p_1p_3,\ p_2p_4p_6$; or $p_5,\ p_1p_6,\ p_2p_3p_4$. The first yields $p_2 \equiv 1 \pmod{p_1}$, $p_1 \equiv 1 \pmod{p_2}$; the second, $p_3 \equiv 1 \pmod{p_1}$, $p_1 \equiv 1 \pmod{p_3}$. For the third case, $S_1 = S_2$ gives

(22) $$p_1p_2=p_3p_4+\delta p_5p_6,\quad \delta+p_1p_3p_4+p_2p_3p_4=p_1p_2p_3+p_1p_2p_4,$$

while $S_1=S_3$ gives $p_5p_6\equiv p_2 \pmod{p_3p_4}$. Thus $\delta\equiv p_1 \pmod{p_3p_4}$, by ($22_1$). Then ($22_2$) gives $p_1\equiv p_1p_2(p_3+p_4)$, mod $p_3p_4$. Hence

(23) $$1=p_2(p_3+p_4)+lp_3p_4 \quad (l \text{ negative integer}).$$

But $p_1p_3p_4p_5p_6\equiv p_3p_4\equiv p_1p_5p_6 \pmod{p_2}$. Thus $p_1p_5p_6\equiv 1$, $p_3p_4\equiv 1 \pmod{p_2}$. Hence in (23), $1\equiv l$, $l=1-mp_2$, $m>0$. Thus (23) may be given the form

$$p_2[(m-1)p_3p_4-p_3-p_4]+p_2p_3p_4=p_3p_4-1.$$

But if $m>1$, the quantity in brackets is positive and the equation impossible. Hence $m=1$, and (23) becomes

(24) $$p_2(p_3p_4-p_3-p_4)=p_3p_4-1.$$

Since $p_2\geqq 2$, the first member exceeds the second if $(p_3-2)(p_4-2)>3$. Since (34) leaves each triple unaltered, we may set $p_3\geqq p_4$. Hence, there remains the cases

$$p_3=5,\ p_4=3;\ p_3=p_4=3;\ p_4=2.$$

For $p_3=5$, $p_4=3$, we have $p_2=2$, $S_2=15+16p_1p_5p_6$, $S_3=30p_5+30p_1p_6+p_1p_5p_6$. Hence $S_2=S_3$ gives $(p_1p_6-2)(p_5-2)=3$, $p_1p_6=5$ or 3, which is impossible. For $p_3=p_4=3$, (24) gives $3p_2=8$. Hence must $p_4=2$, $p_2(p_3-2)=2p_3-1$. Thus $p_2\geqq 3$, and the first member exceeds the second if $p_3>5$. Hence $p_3=5$, $p_2=3$, or $p_3=3$, $p_2=5$. For the first, $2S_2=2S_3$ becomes $(2p_5-3)(2p_1p_6-3)=7$, whence $p_1p_6=5$ or 2. For the second, $4S_2=4S_3$ becomes $(4p_1p_6-5)(4p_5-5)=21$, whence $p_1p_6=3$ or 2.

Next, let two of the triples be the first two in (21) and the third of type (2, 2, 2). The latter may be assumed to be $p_1p_2$, $p_3p_4$, $p_5p_6$; $p_1p_2$, $p_3p_5$, $p_4p_6$; $p_1p_3$, $p_2p_4$, $p_5p_6$; or $p_1p_3$, $p_2p_5$, $p_4p_6$. The first two and the last two are excluded by the argument excluding the first and second cases, respectively, at the beginning of the preceding paragraph.

*Hence at most one of the three triples is of the type* (1, 1, 4).

9. Let two triples be $p_1$, $p_2$, $p_3p_4p_5p_6$ and $p_3$, $p_1p_2$, $p_4p_5p_6$. Adding $p_1\ldots p_6$ to $S_1$ and $S_2$ and equating the sums, we get

(25) $$p_1p_2(p_3-1)(p_4p_5p_6-1)=p_3p_4p_5p_6(p_1-1)(p_2-1).$$

We may set $p_2 \geqq p_1$, $p_2 \neq p_3$. Hence $p_2$ divides $p_4p_5p_6$, so that we may set $p_4=p_2$. Since $p_4p_5p_6-1>p_5p_6(p_2-1)$, we have $p_1(p_3-1)<p_3(p_1-1)$, $p_1>p_3$. Hence $p_2>p_3$. If the third triple is of type (1, 2, 3), it may be taken to be $p_5$, $p_1p_3$, $p_2^2p_6$; $p_5$, $p_1p_6$, $p_2^2p_3$; $p_5$, $p_2p_3$, $p_1p_2p_6$; $p_5$, $p_2^2$, $p_1p_3p_6$; $p_5$, $p_2p_6$, $p_1p_2p_3$; or $p_5$, $p_3p_6$, $p_1p_2^2$. Now $S_1$ is a multiple of $p_2$, so that $S_3$ must be. Hence for the first case, $p_1p_3p_5$ is a multiple of $p_2$, whence $p_1=p_2$. Then $S_1$ is a multiple of $p_2^2$, so that $S_3$ must be; hence $p_3p_5\equiv 0 \pmod{p_2}$, which is impossible. For the second case, $p_1p_5p_6\equiv 0$ $\pmod{p_2}$. If $p_1=p_2$, $S_1\equiv 0 \pmod{p_2^2}$, so that by $S_3$, $p_6=p_2$. In any event, $p_6=p_2$ and the second and third triples have $p_1p_2=p_1p_6$ in common. For the third case, $S_1=S_2$ gives $p_1p_3=p_1+^{\varepsilon}p_5p_6$, whence $^{\varepsilon}=^{\delta}p_1$; while $S_1=S_3$ gives $p_1p_2\equiv p_1p_2p_5p_6$, $p_5p_6\equiv 1 \pmod{p_3}$, $p_5p_6>p_3$. For the fourth case, $S_1\equiv S_3$ mod $p_2$ gives $p_1p_3p_6p_5\equiv 0 \pmod{p_2}$. If $p_1=p_2$ the second and third triples would have a common element. Hence $p_2=p_6$. By (25), $p_5p_6$ must divide $p_1(p_3-1)$. But $p_2>p_3-1$. Hence $p_2=p_6$ requires $p_1=p_2$. For the fifth case, $p_1\neq p_6$ by $p_2p_6$ and $p_1p_2$. Removing the factor $p_2$ from the $S$'s, we see that $p_2p_3p_5p_6\equiv p_3p_5p_6\equiv p_5p_6 \pmod{p_1}$, $p_3\equiv 1 \pmod{p_1}$, contrary to $p_3<p_1$. For the sixth case, $S_1\equiv S_3 \pmod{p_2}$ gives $p_3p_5p_6\equiv 0$ $\pmod{p_2}$, $p_6=p_2$. As in the fourth case, $p_1=p_2$. Then $S_1$, but not $S_3$, is a multiple of $p_2^2$.

If the third triple is of type (2, 2, 2), it may be taken to be

$$p_2^2,\ p_1p_3,\ p_5p_6;\ p_2^2,\ p_1p_5,\ p_3p_6;\ p_2p_3,\ p_2p_5,\ p_1p_6;\ \text{or}\ p_2p_5,\ p_2p_6,\ p_1p_3.$$

For the first two cases, $p_1\neq p_2$ and $S_1\equiv S_3 \pmod{p_2}$ gives $p_6=p_2$, contrary to the above. The third case is excluded by $p_3\equiv 1 \pmod{p_5p_6}$, $p_5p_6\equiv 1$ $\pmod{p_3}$; the fourth by $p_1\equiv 1 \pmod{p_3}$, $p_3\equiv 1 \pmod{p_1}$.

THE UNIVERSITY OF CHICAGO, *May 16, 1909.*

◆

# NOTE ON SOME POLYNOMIALS RELATED TO LEGENDRE'S COEFFICIENTS.*

By **R. D. CARMICHAEL, Anniston, Alabama.**

The object of this note is to point out some properties of a class of functions which contains Legendre's coefficients as a special case. It will be seen that the former possess some interesting properties belonging to the latter.

1. Consider the definite integral†

---

*Presented to the American Mathematical Society, April 24, 1909.

†Cf. Goursat-Hedrick, *Mathematical Analysis*, Vol. I, p. 173.